

MACRO-INVERTEBRATE COMMUNITIES OF BAA ATOLL, REPUBLIC OF MALDIVES

BY

SERGE ANDRÉFOUËT,¹ JEAN-LOUIS MENOU,¹ AND SHAFIYA NAEEM²

ABSTRACT

In order to guide conservation efforts in Baa Atoll in the Republic of Maldives, macro-invertebrate species (principally sponges, sea stars, urchins, holothurians, crinoids, ophiuroids, ascidians, bivalves, gastropods, anemones, zooanthids, anthipatharian, gorgonians, alcyonaceans, and flatworms) were inventoried on 21 sites to provide an index of benthic species richness (number of species). A total of 182 species were recorded and identified. Richness ranged between 6 to 40 species per sampling sites. The richest sites were hard-bottom submerged lagoonal patch reefs experiencing high currents. Ecological rarity dominates the macro-invertebrate community pattern with 95 species found in only one sampling site. This new data set brings fresh knowledge on Baa Atoll and Maldives in order to identify biodiversity hot-spots. Further work will investigate the sensitivity of biodiversity conservation planning and sitting algorithms to this data set.

INTRODUCTION

Invertebrates include an extremely wide range of organisms with 8 major phyla (Porifera, Cnidaria, Annelida, Sipuncula, Arthropoda, Mollusca, Echinodermata, Chordata), and 24 minor phyla (see Glynn and Enochs, (2011) for a complete list). Coral reefs offer the richest biodiversity on Earth, along with tropical rainforest (Reaka-Kudla, 1997), and the greatest part of this diversity lies in the invertebrate communities. Glynn and Enochs (2011) recently reviewed the taxonomy and functional roles of invertebrates in coral reefs. They offer a synoptic view of the complexity of the invertebrate world. Here, we will consider more particularly the macro-invertebrates. The definition of the macro-invertebrate community is fuzzy, and context-dependant. For instance, Glynn and Enochs (2011) distinguished for cryptic species the macrofauna from the meiofauna, depending if the specimen body size is larger than 1 mm or not. Thus, body sizes, associated with investigation means and habitats define in a relative way the community of macro-invertebrates. Here, the macro-invertebrate benthic community includes all species that can be detected by eye while diving or snorkeling, on soft to hard-bottoms, on both biotic and abiotic substrates.

¹ Institut de Recherche Pour le Développement, RD Nouméa, UMR 227 COREUS, BP A5, 98800 Nouméa cedex, New-Caledonia

² Marine Research Center, H. White waves, Moonlight Higun, Malé – 20025, Maldives

Trying to measure invertebrate biodiversity for any given reef complex such as a Maldivian atoll is a daunting challenge that requires enormous field work efforts (Bouchet et al., 2011), rare taxonomic expertise, and genetic analyses. Thus, coral reef sites compendium providing within an accepted taxonomic reference the full range of existing species for most of the 32 phyla are extremely rare (e.g., New-Caledonia in Payri and Richer de Forges, 2006; Guam in Paulay 2003, Santo Island, Vanuatu in Bouchet et al. 2011).

Among valuable resources, a number of specialized taxonomic publications list macro-invertebrate species for selected phyla and locations (e.g., for New Caledonia: echinoderms, ascidians, sponges, gorgonians, are described respectively in Guille et al. 1986, Moniot et al., 1991; Levi et al. 1998; Grasshoff and Bargibant, 2001) but no recent taxonomic compilation exists for Maldives macro-invertebrates to our knowledge. The previous reference was Clarke and Rowe (1971) which is now outdated for several species.

Several well illustrated field guide books list invertebrates species for a given region (e.g., Colin and Arneson, 1995; Coleman, 2001; Laboute and Richer de Forges, 2004; Hervé, 2010; Poupin and Juncker, 2010). Web-based databases and information systems are also increasingly set to catalogue and identify species, and infer distribution maps (e.g., the integrated Taxonomic Information System, <http://www.itis.gov>, the World Register of Marine Species <http://www.marinespecies.org/index.php>, the Ocean Biogeographic Information System, <http://v2.iobis.org/> and its Indian Ocean node <http://www.indobis.org/>). Nevertheless, to measure and characterize the biodiversity of a particular previously undocumented coral reef site, often, only a limited number of phyla (e.g., Cnidaria), classes (e.g., Anthozoa), orders (e.g., Scleractinia) are targeted for new collections. Even more often, in a conservation or monitoring context, census and characterization focus only on few indicator species from various phyla, typically commercial and functionally prominent species of mollusk, crustaceans and echinoderms. This limited approach, taxonomically speaking, corresponds to rapid assessments as performed for instance by NGOs (e.g., Conservation International, McKenna et al., 2009; The Nature Conservancy, Ramohia, 2006), although some taxa have been more thoroughly investigated (e.g., mollusks in McKenna et al., 2002, Appendix 3).

To provide a measurement of Baa Atoll invertebrate biodiversity as completed as possible given the logistic and expertise constraints, it was decided to provide for the most dominant atoll habitats a list of conspicuous macro benthic invertebrates found during day time only. The rationale is that this macro-fauna assessment would provide an index of biodiversity to characterize reef habitat benthic richness and to guide the identification of biodiversity hot-spots for conservation planning (Hamel and Andréfouët, this issue). We report here on the variation of this richness in Baa Atoll. Therefore, the invertebrate sampling effort performed in Baa Atoll can not be compared to huge efforts such as those conducted recently in Santo Island for instance (Bouchet et al., 2011), but it provides an enhanced view of Baa Atoll communities richer than most typical rapid assessment programs that focussed on commercial species only (Le Berre et al., 2009). The Baa census included scleractinian (hard corals) and hydrozoans. These taxa are treated elsewhere (Bigot and Hamir, this issue; Gravier-Bonnet and Bourmaud, this issue). The present census did not consider crustaceans for time and logistical reasons.

METHODOLOGY

Macro-invertebrates communities were sampled during day dives conducted between 26 May to 4 June 2009. During each dive, specimen were inventoried from the deeper areas (typically around 40 m) to the shallows following a random path. This technique is generally more rewarding for biodiversity census than working through transects and quadrats that limit the searching spatial domain. Specimens were photographed in situ, and brought on board the diving vessel for further examination and identification when needed. Specimen were identified at the highest possible taxonomic level (species or genera). No museum collections were created.

Sampling sites were selected to cover the dominant reef types and exposure in Baa Atoll. This included outer oceanic slope and flats, submerged lagoon patch reefs, and sheltered central lagoonal slopes and reef flats. A total of 21 sites were sampled across Baa Atoll (Fig. 1).

Richness per station was simply defined as the number of species, all phyla included, found during the survey.

RESULTS AND DISCUSSION

The survey of the 21 stations provided a total of 182 identified species (Table 1 in Appendix), plus 5 specimen of sponges and 2 Crinoidea species left unidentified. Records were distributed among the phylum Cnidaria (class Anthozoa, $n=34$), Echinodermata (class Asteroidea $n=10$, Crinoidea $n=9$, Echinoidea $n=7$, Holothuroidea $n=12$, Ophiuroidea $n=11$), Mollusca (class Bivalvia $n=14$, Cephalopoda $n=1$ and Gastropoda $n=48$), Porifera (class Calcarea $n=3$, Demospongiae $n=26$), Chordata (class Ascidiacea, $n=7$), Annelida (class Polychaeta, $n=1$) and Platyhelminthes (class Turbellaria, $n=1$). Selected specimen are illustrated Plates 1 to 5.

Richness per station ranged between 6 (station 23: deep lagoon, and station 2: seagrass bed) to 40 species (station 1, lagoon slope and reef flat). Figure 2 and Figure 3 provide the continuum of richness found for all stations. In terms of richness per habitat, the submerged lagoon patch reef frequently provided high richness ($N>29$) with stations 10, 11, 17, 25. These stations were characterized by high currents, hard-bottom and numerous hang-outs and small caves. Station 19, one of the poorest stations with 12 species, was also a patch reef characterized by high current but also by soft bottom, in contrast with the richer patch reef sites.

Average richness per geomorphological habitat type varied (Fig. 4), but the poorest habitats (deep lagoon, oceanic reef flat, and seagrass beds) where also the less sampled ($n=1$ for each of them). For the well sampled habitats ($n>5$), richness appeared homogeneous without significant difference (Fig. 4). The survey that occurred only on lagoon reef flats (thus without adjacent deep slopes) yielded a similar richness ($N=26$, $n=1$, station 3) than the dives that occurred on both lagoon slopes and flats ($N=23.5$, $n=7$), even if they have been visited only once. Thus, a sampling effort bias on lagoon reef flats was likely less an issue to estimate richness, compared to deep lagoons and outer reef flats.




Figure 1. Location of Baa Atoll and macro-invertebrates sampling stations.

The converging richness values for different strata suggest that around 35 species per habitat is the richness limit that can be reached given the sampling method (Fig. 3 and Fig. 4). Obviously, additional species will appear by searching in cryptic spaces, by night, during longer dives, and for more sampling sites per strata.

The turn over of species per station was high, highlighting ecological rarity in Baa Atoll macro-invertebrates community. More than half of the species ($n=95$) were found in only one station, suggesting that macro-invertebrates beta-diversity as it was sampled here in Baa Atoll reefs is high (Fig. 4). The most frequently encountered species were the starfish *Linckia multifora* (found on 17 stations), the pin-cushion seastar *Culcita schmedeliana* (15 stations) and the holothurian *Pearsonothuria graeffei* (15 stations). Two specimen of *Acanthaster planci* were found in station 25 only during the macro-invertebrate surveys, although others have been seen during the habitat survey in station 23.




Figure 2. Variation of species richness per sampled station.

Similar high turnover of species was noted for macromollusc species in Tuamotu atolls in the Central Pacific Ocean (Pante et al., 2006), with 70% of 27 species encountered in only one or two habitats. A systematic comparison with the study by Adjéroud et al. (2000) and Pante et al. (2006) is not possible since they focused on lagoon habitats only and considered different type of habitats and geomorphological strata than the present study, and not all the phylum considered here. However, as seen in Figure 4, lagoon habitats sheltered an average of about 25 species in Baa Atoll. All lagoon habitats yielded a total of 94 species mollusks and echinoderms species. In average, lagoon stations provided 15.0 ± 3.9 species (mean \pm stdev, $n=13$). In contrast, Adjéroud et al. (2000) listed a grand total of 36 species in 9 atolls, thus nearly 1/3 less than in Baa alone. They also reported a range of lagoon-scale richness between 2 and 22 for 9 atoll lagoons, with 10.4 ± 7.1 species (mean \pm stdev, $n=9$).

Pearse (2009) reported for shallow water asteroids, echinoids and holothuroids in 6 Pacific Ocean sites that 56% of the 113 species were found in only one site. Thus, we found a similar ratio of single species in Baa Atoll than in a Pacific wide search suggesting that rarity could obey pattern across multiple spatial and biogeographical scales but this warrants larger tropical investigations on invertebrates community structure (Iken et al., 2010).




Figure 3. Mapped species richness for the 21 stations (Fig. 1).




Figure 4. Variation of richness per habitat.




Figure 5. Histogram of species occurrences for the 21 sites.

Similar observations on high beta-diversity can be made at the genera level. 127 genera were identified and recorded, and 50 were found only once. Half of the genera were found only once or twice. The most frequently found genera were the echinoderms *Linckia*, *Culcita*, *Pearsonothuria* and *Choriaster* found in 18, 17, 15, and 13 stations respectively. Next was the mollusk genera *Tridacna* with 11 stations.

Few recent (in the past 20 years) publications have updated taxonomic check-lists for macro-invertebrates in the Maldives (e.g. for 47 species of brachyuran crabs in Laamu Atoll, see Kumar and Wesley, 2010) and none to our knowledge which would be Baa-specific. Commercial macro-invertebrate resources were the object of national-scale fishery reports, specifically for clams (Basker, 1991) and sea cucumbers (Joseph, 1992). Both authors reported that these fisheries were already unsustainable with quickly depleting resources. Clams (both *Tridacna maxima* and *T. squamosa*) were actually more present in Baa than what Table 1 suggests. Indeed, they were seen on virtually all stations of the survey completed for habitat mapping (Andréfouët et al., this issue). They may have been missed during the macro-invertebrate survey given the longer time spent at depth than in the shallows (between 0 and 5 meters) where they are more abundant. For sea cucumbers, Joseph (1992) listed 9 species commonly fished throughout Maldives, and refer to an 1988 expedition by a Maldives/China UNDP/ESCAP pilot study that listed 8 species identified from Baa, Haa Alifu and Haa Dhaalu Atolls, namely *Bohadschia marmorata*, *Actinopyga lecanora*, *A. mauritiana*, *Holothuria (Halodeima) atra*, *H. leucospilota*, *H. (Microthele) nobilis*, *Stichopus chloronotus*, and *Synapta maculata*. The last four of these 8 species were not found in the 2009 Baa survey which yielded a total of 12 species (Table 1).

In Diego Garcia, an atoll of the Chagos Archipelago in the south of Maldives, Clark and Taylor (1971) reported for the Echinodermata phyla, 9, 11, 1 and 14 species for respectively the Echinoidea, Ophiuroidea, Asteroidea and Holothuroidea classes. In Baa, we report here respectively 7, 11, 10 and 12 species for the same classes, thus similar values, except for the Asteroidea. Conversely, Taylor (1971) reported for Diego Garcia 179 species of mollusks alone, *vs* 63 species here (class Bivalvia $n=14$, Cephalopoda $n=1$ and Gastropoda $n=48$). Overall, the Mollusca phylum appears undersampled in Baa compared to other check-lists published for the region in the past (e.g. Kohn and Robertson (1968) for conidae).

CONCLUSION

The image of the benthic fauna made from the distribution of sponges, sea stars, urchins, holothurians, crinoids, ophiuroids, ascidians, bivalves, gastropods, anemones, zooanthids, anthipatharian, gorgonians, alcyonaceans, and flatworms provided a fresh view of the Baa Atoll benthic macro-invertebrate richness. This assessment is obviously dependent on the sampling strategy applied in May-June 2009, and for the sake of knowledge more detailed census with larger resources remain worth investigating for Baa, and for the Maldives in general. Nevertheless, the results presented here update the level of knowledge of Baa atoll and Maldives biodiversity.

The taxonomic inventory performed here brings a different biodiversity layer to guide conservation planning than what was usually done for coral reef ecosystems. Conservation planning is dependent on optimization algorithms, and these algorithms often use richness and rarity, and complementarity between sites to converge towards a solution able to fill the conservation targets. Given the beta-diversity patterns observed here, further work will investigate the sensitivity of conservation planning and siting algorithms to the macro-invertebrate layer.

ACKNOWLEDGEMENTS

This work is a contribution to the program “Biodiversity of Indian Ocean Islands” of the French Fondation pour la Recherche sur la Biodiversité. Gratitude is expressed to Shiham Adam, Hussain Zahir and Thomas Le Berre for scientific collaboration during the field survey, and to the Atoll Ecosystem Conservation Project for funding.

REFERENCES

- Adjeroud, M., S. Andréfouët, C. Payri, and J. Orempuller.
2000. Physical factors of differentiation in macrobenthic communities between atoll lagoons in the Central Tuamotu Archipelago (French Polynesia). *Marine Ecology Progress Series* 196:39-48.
- Basker, J.
1991. Giant clams in the Maldives. A stock assessment and study of their potential for culture. Bay of Bengal Program. Food and Agriculture Organisation of the United Nations, Report BOPB/WP/72, Madras, India. 24 p.
- Bouchet, P., Le Guyader, H. and Pascal, O.
2011. The natural history of Santo. Vol. 69. Muséum National d'Histoire Naturelle, Paris ; Institut de Recherche pour le Développement, Paris, Marseille ; Pro-Natura International, Paris. 572 p.
- Clark, A.M., and F. W. E. Rowe
1971. Monograph of shallow water Indo-Pacific echinoderms. British Museum (Natural History), London, Pub. 690. 306 p. (with Plates and references)
- Clark, A. M., and J. D. Taylor
1971. Echinoderms from Diego Garcia. *Atoll Research Bulletin* 149:89-92
- Taylor, J.D.
1971. Marine Mollusca from Diego Garcia. *Atoll Research Bulletin* 149:105-126
- Coleman, N.
2000. Marine life of the Maldives. Atoll Editions. 352 p.
- Colin, P., and C. Arneson
1995. Tropical Pacific Invertebrates: a field guide to the marine invertebrates occurring on tropical Pacific coral reefs, seagrass beds, and mangroves. Coral Reef Press, Beverly Hills, USA. 296 p.
- Glynn, P. W., and I. C. Enochs
2011. Invertebrates and their roles in coral reef ecosystems. Pages 273-325 in Z. Dubinsky and N. Stambler (Eds) *Coral reefs: an ecosystem in transition*. Springer, Berlin.
- Guille, A., P. Laboute, and J. Menou
1986. Handbok of the sea-stars, sea-urchins, and related echinoderms of New-Caledonia lagoon. ORSTOM, Paris. 239 p.
- Grasshoff, M., and G. Bargibant
2001. Coral reef gorgonians of New Caledonia. IRD, Paris. 336 p.
- Hervé, J.F
2010. Guides des nudibranches de Nouvelle-Calédonie et autres Opisthobranches. Ed. Catherine Ledru – IRD, Nouméa, Nouvelle-Calédonie, 401 p.
- Iken, K. and 17 authors
2010. Large-scale spatial distribution patterns of echinoderms in nearshore rocky habitats. *Plos ONE* 5 :e13845
- Joseph, L.
1992. Review of the bêche de mer (sea cucumber) fishery in the Maldives. Bay of Bengal Program. Food and Agriculture Organisation of the United Nations, report BOPP/WP/79 Madras, India. 34 p.

- Kohn, A. J., and R. Robertson.
1968. The Conidae (Gastropoda) of the Maldives and Chagos Archipelagoes. *J. Mar. Biol. Ass. India* 8: 273-277
- Kumar, A. A. J., and S. G. Wesley.
2010. Spatio-temporal distribution of assemblages of brachyuran crabs at Laamu Atoll, Maldives. *Atoll Research Bulletin* 588:1-15.
- Laboute, P., and B. Richer de Forges
2004. Lagons et récifs de Nouvelle-Calédonie. Editions Ledru, IRD, Nouméa. 520 p.
- Le Berre T., Emanuelli E., Roussel E., Guignard C., Sivad A., Saleem M., Zahir H.
2009. Rapid marine ecological baseline assessment of islands of Baa atoll. *Technical report SEAMARC, Atoll Ecosystem Conservation Project*, 156 pp.
- Lévi, C., P. Laboute, G. Bargibant, and J. Menou
1998. Sponges of the New Caledonia lagoon. ORSTOM, Paris. 215 p.
- McKenna, S. A., G. R. Allen, and S. Suryadi.
2002. A marine rapid assessment of the Raja Ampat Islands, Papua Province, Indonesia. RAP Bulletin 22. Conservation International, Washington. 195 p.
- McKenna, S. A., N. Baillon, and J. Spaggiari.
2009. Rapid marine biodiversity assessment of the coral reefs of the northwest lagoon, between Koumac and Yandé, Province Nord, New Caledonia. RAP Bulletin of Biological Assessment 53. Conservation International, Washington. 184 p.
- Monniot, C., F. Monniot, and P. Laboute
1991. Coral reef ascidians of New Caledonia. ORSTOM, Paris. 248 p.
- Pante, E., M. Adjeroud, P. Dustan, L. Penin, and M. Schrimm.
2006. Spatial patterns of benthic invertebrate assemblages within atoll lagoons: importance of habitat heterogeneity and considerations for marine protected area design in French Polynesia. *Aquatic Living Resource* 19:207-217.
- Paulay, G. H.
2003. Marine biodiversity of Guam and the Mariannas. *Micronesica*. 35-36:456-513
- Payri, C., and B. Richer de Forges
2007. Compendium of marine species of New Caledonia. IRD, Centre de Nouméa, Nouméa. 435p. + 19 plates
- Pearse, J.
2009. Shallow water asteroids, echinoids and holothuroids at 6 sites across the tropical west Pacific; 1988-1989. *Galaxea* 11:187-195
- Poupin, J. and M. Juncker
2010. A guide to the decapode crustacean of the South Pacific/ Guide des crustacés décapodes du Pacifique Sud. CRISP/SPC, Nouméa, Nouvelle-Calédonie, 320 p.
- Ramohia, P.
2006. Fisheries resources: commercially important macroinvertebrates. In: Green, A., P. Lokani, W. Atu, P. Ramohia, P. Thomas and J. Almany (eds). 2006. Solomon Islands Marine Assessment: Technical report of survey conducted May 13 to June 17, 2004. TNC Pacific Island Countries Report No 1/06. pp.329-400

Table 1. Check list of macro-invertebrates species for the 21 sampled stations (1=presence).

Phylum	Class	Order	Genera	Species	1	2	9	4	9	7	9	10	11	10	17	19	21	28	28	25	27	28
Annelida	Polychaeta	Sabellida	Sabellastarte	indica	1													1				
Chordata	Asciacea	Aplosobranchia	Didemnum	molle	1	1								1	1	1				1	1	1
Chordata	Asciacea	Aplosobranchia	Diplosoma	versicolor													1					
Chordata	Asciacea	Aplosobranchia	Eudistoma	sp.																	1	
Chordata	Asciacea	Aplosobranchia	Polysynchraton	sp.		1									1							
Chordata	Asciacea	Plebobranchia	Ascidia	glabra									1	1	1							
Chordata	Asciacea	Stolidobranchia	Botrylloides	sp.									1									
Chordata	Asciacea	Stolidobranchia	Pyura	gangelion		1								1								
Cnidaria	Anthozoa	Actinaria	Actinodendron	arborum	1														1	1		
Cnidaria	Anthozoa	Actinaria	Heteractis	magnifica					1			1										
Cnidaria	Anthozoa	Actinaria	Stichodactyla	cf mertensii					1			1										
Cnidaria	Anthozoa	Actinaria	Doffeina	sp.								1				1						
Cnidaria	Anthozoa	Actinaria	Heteractis	aurora								1										
Cnidaria	Anthozoa	Actinaria	Actinodendron	cf. plumosum														1				
Cnidaria	Anthozoa	Actinaria	Discosoma	sp.														1				
Cnidaria	Anthozoa	Actinaria	Entacmea	quadricolor															1	1		
Cnidaria	Anthozoa	Alcyonacea	Dendronephthya	sp. 1						1		1			1							
Cnidaria	Anthozoa	Alcyonacea	Chironephthya	sp. 1								1			1							
Cnidaria	Anthozoa	Alcyonacea	Chironephthya	sp. 2											1							
Cnidaria	Anthozoa	Alcyonacea	Dendronephthya	sp. 2							1											
Cnidaria	Anthozoa	Alcyonacea	Dendronephthya	sp. 3									1									
Cnidaria	Anthozoa	Alcyonacea	Subergorgia	sp.							1											
Cnidaria	Anthozoa	Alcyonacea	Acanthogorgia	sp. 1						1												
Cnidaria	Anthozoa	Alcyonacea	Acabaria	sp. 1				1				1			1							
Cnidaria	Anthozoa	Alcyonacea	Acabaria	sp. 2					1													
Cnidaria	Anthozoa	Alcyonacea	Acabaria	sp. 3								1			1							
Cnidaria	Anthozoa	Alcyonacea	Annella	mollis												1						
Cnidaria	Anthozoa	Alcyonacea	Acanthogorgia	sp. 2											1							
Cnidaria	Anthozoa	Alcyonacea	Acanthogorgia	sp. 3										1								
Cnidaria	Anthozoa	Antipatharia	Stichopathes	sp.	1							1	1	1	1							
Cnidaria	Anthozoa	Antipatharia	Cirripathes	sp. cf. anguineus	1							1		1								
Cnidaria	Anthozoa	Antipatharia	Antipathes	sp. 1						1			1									
Cnidaria	Anthozoa	Antipatharia	Antipathes	sp. 3				1							1							
Cnidaria	Anthozoa	Antipatharia	Antipathes	sp. 4											1							
Cnidaria	Anthozoa	Antipatharia	Antipathes	sp. 5								1		1	1							
Cnidaria	Anthozoa	Antipatharia	Antipathes	sp. 6											1							
Cnidaria	Anthozoa	Zoantheidea	Palythoa	caesia	1	1	1	1				1									1	
Cnidaria	Anthozoa	Zoantheidea	Zoanthus	mantoni															1			
Cnidaria	Anthozoa	Zoantheidea	Protopalythoa	sp.																1		1
Cnidaria	Anthozoa	Zoantheidea	Zoanthus	sp.					1													
Cnidaria	Anthozoa	Zoantheidea	Parazoanthus	sp.											1							
Cnidaria	Anthozoa	Zoantheidea	Palythoa	caesia cf. tuberculosa																		
Echinodermata	Astroidea	Spinulosida	Acanthaster	planci																		
Echinodermata	Astroidea	Valvatida	Choriaster	granulatus	1	1	1	1				1	1	1	1		1		1	1	1	1
Echinodermata	Astroidea	Valvatida	Culcita	schmedeliana	1	1	1	1				1	1	1	1		1		1	1	1	1
Echinodermata	Astroidea	Valvatida	Linckia	guidingi	1				1	1											1	1
Echinodermata	Astroidea	Valvatida	Culcita	cf. novaeguineae					1									1				
Echinodermata	Astroidea	Valvatida	Linckia	multiflora	1	1	1	1				1	1	1	1	1	1	1	1	1	1	1
Echinodermata	Astroidea	Valvatida	Fromia	indica			1										1					
Echinodermata	Astroidea	Valvatida	Gomophia	sp.														1				
Echinodermata	Astroidea	Valvatida	Fromia	milleporella										1								
Echinodermata	Astroidea	Valvatida	Fromia	nodosa								1						1	1	1		
Echinodermata	Astroidea	Valvatida	Comanthus	parvicirrus				1						1	1	1						
Echinodermata	Crinoidea	Comatulida	Comanthina	schlegeli		1						1	1									

Table 1 (Con'td)

Phylum			Class			Order			Genera			Species			Stations													
1	2	3	4	5	5	7	5	10	11	10	17	10	21	22	23	23	25	27	28									
Echinodermata	Crinoidea	Comatulida	Capillaster	sp.				1	1	1			1				1											
Echinodermata	Crinoidea	Comatulida	Comanthus	sp. 2							1																	
Echinodermata	Crinoidea	Comatulida	Comaster	sp.								1		1														
Echinodermata	Crinoidea	Comatulida	Comanthus	cf. parvicirrus							1						1											
Echinodermata	Crinoidea	Comatulida	Stephanometra	sp.										1														
Echinodermata	Echinoidea	Diadematoidea	Echinothrix	calamaris			1																					
Echinodermata	Echinoidea	Diadematoidea	Diadema	savignyi																								
Echinodermata	Echinoidea	Echinoidea	Echinostrephus	molaris							1																	
Echinodermata	Echinoidea	Echinoidea	Heterocentrotus	mammillatus																								
Echinodermata	Echinoidea	Echinoidea	Echinometra	mathaei																								
Echinodermata	Echinoidea	Echinoidea	Echinometra	sp1-cf sternalis																								
Echinodermata	Echinoidea	Spatangoida	Metalia	sp.			1							1														
Echinodermata	Echinoidea	Spatangoida	Schizaster	sp.								1																
Echinodermata	Holothuroidea	Apodida	Synaptula	sp.																								
Echinodermata	Holothuroidea	Aspidochirotida	Pearsonothuria	graeffe																								
Echinodermata	Holothuroidea	Aspidochirotida	Holothuria	fuscogilva																								
Echinodermata	Holothuroidea	Aspidochirotida	Actinopyga	lecanora																								
Echinodermata	Holothuroidea	Aspidochirotida	Bohadschia	argus																								
Echinodermata	Holothuroidea	Aspidochirotida	Bohadschia	marmorata sp1																								
Echinodermata	Holothuroidea	Aspidochirotida	Bohadschia	marmorata sp2																								
Echinodermata	Holothuroidea	Aspidochirotida	Actinopyga	mauritiana																								
Echinodermata	Holothuroidea	Aspidochirotida	Thelenota	ananas																								
Echinodermata	Holothuroidea	Aspidochirotida	Thelenota	anax																								
Echinodermata	Holothuroidea	Aspidochirotida	Holothuria (Halodeima)	edulis																								
Echinodermata	Holothuroidea	Aspidochirotida	Holothuria (Halodeima)	atra																								
Echinodermata	Holothuroidea	Aspidochirotida	Holothuria (Acanthophiothrix)	sp1																								
Echinodermata	Ophiuroidea	Ophiurida	Ophiothrix	danae																								
Echinodermata	Ophiuroidea	Ophiurida	Ophiothela	erinaceus																								
Echinodermata	Ophiuroidea	Ophiurida	Ophiocoma	cf. cincta																								
Echinodermata	Ophiuroidea	Ophiurida	Ophiolepis	elegans																								
Echinodermata	Ophiuroidea	Ophiurida	Ophiarthrum	sp. 2																								
Echinodermata	Ophiuroidea	Ophiurida	Ophiothrix	sp. 3																								
Echinodermata	Ophiuroidea	Ophiurida	Ophionereis	sp. 1																								
Echinodermata	Ophiuroidea	Ophiurida	Ophiarachnella	cf. gorgonia																								
Echinodermata	Ophiuroidea	Ophiurida	Ophiothrix (Acanthophiothrix)	purpurea																								
Echinodermata	Ophiuroidea	Ophiurida	Ophionereis	sp. 2																								
Mollusca	Bivalvia	Arcoida	Arca	ventricosa																								
Mollusca	Bivalvia	Ostreoida	Hyotissa	hyotis																								
Mollusca	Bivalvia	Ostreoida	Lopha	cristagalli																								
Mollusca	Bivalvia	Pectinoida	Spondylus	varians																								
Mollusca	Bivalvia	Pectinoida	Spondylus	sp.																								
Mollusca	Bivalvia	Pterioidea	Pinctada	margaritifera																								
Mollusca	Bivalvia	Pterioidea	Pinna	sp.																								
Mollusca	Bivalvia	Pterioidea	Pteria	penguin																								
Mollusca	Bivalvia	Veneroidea	Cardiidae	indéterminé																								
Mollusca	Bivalvia	Veneroidea	Cardium	sp.																								
Mollusca	Bivalvia	Veneroidea	Globivenus	toreuma																								
Mollusca	Bivalvia	Veneroidea	Tapes	literatus																								
Mollusca	Bivalvia	Veneroidea	Tridacna	maxima																								
Mollusca	Bivalvia	Veneroidea	Tridacna	squamosa																								
Mollusca	Bivalvia	Veneroidea	Octopus	cyanea																								
Mollusca	Cephalopoda	Octopoda	Cerithium	nodulosum																								
Mollusca	Gastropoda	Caenogastropoda	Rhinoclavis	vertagus cf.																								
Mollusca	Gastropoda	Caenogastropoda	Rhinoclavis	aspera																								
Mollusca	Gastropoda	Littorinimorpha	Cypraea	tigris																								

Table 1 (Con'td)

			Stations																				
Phylum	Class	Order	Genera	Species	1	2	3	4	5	5	7	10	11	10	17	19	10	21	23	23	25	27	23
Mollusca	Gastropoda	Littorinimorpha	Cypraea	caputserpentis	1						1												
Mollusca	Gastropoda	Littorinimorpha	Cypraea	tigris																			1
Mollusca	Gastropoda	Littorinimorpha	Cypraea	linx							1												
Mollusca	Gastropoda	Littorinimorpha	Dendropoma	maximum							1												
Mollusca	Gastropoda	Littorinimorpha	Lambis	lambis																			1
Mollusca	Gastropoda	Littorinimorpha	Lambis	chiragra arthritica								1									1		
Mollusca	Gastropoda	Littorinimorpha	Lambis	truncata truncata							1										1		
Mollusca	Gastropoda	Littorinimorpha	Monoplex	aquatilis																			1
Mollusca	Gastropoda	Littorinimorpha	Septa	rubecula							1												
Mollusca	Gastropoda	Littorinimorpha	Serpulorbis	grandis	1		1		1		1				1				1				
Mollusca	Gastropoda	Littorinimorpha	Strombus (canarium)	sp.cf. dentatum	1										1								
Mollusca	Gastropoda	Neogastropoda	Acus	crenulatus																			
Mollusca	Gastropoda	Neogastropoda	Benimakia	nodata																1			
Mollusca	Gastropoda	Neogastropoda	Chicoreus	microphyllus	1										1								
Mollusca	Gastropoda	Neogastropoda	Chicoreus	sp2				1				1			1								
Mollusca	Gastropoda	Neogastropoda	Chicoreus	palmarosae	1						1											1	
Mollusca	Gastropoda	Neogastropoda	Chicoreus	ramosus																			
Mollusca	Gastropoda	Neogastropoda	Conus	eburneus			1																
Mollusca	Gastropoda	Neogastropoda	Conus	lithoglyphus											1								
Mollusca	Gastropoda	Neogastropoda	Conus	virgo	1																		
Mollusca	Gastropoda	Neogastropoda	Conus	miliaris			1																
Mollusca	Gastropoda	Neogastropoda	Conus	miles cf.												1							1
Mollusca	Gastropoda	Neogastropoda	Conus	arenatus cf.																			1
Mollusca	Gastropoda	Neogastropoda	Conus	emaciatius cf.								1											1
Mollusca	Gastropoda	Neogastropoda	Conus	bandanus																			
Mollusca	Gastropoda	Neogastropoda	Conus	distans							1												
Mollusca	Gastropoda	Neogastropoda	Drupa	sp.																			1
Mollusca	Gastropoda	Neogastropoda	Drupa	ricina arachnoides	1						1												
Mollusca	Gastropoda	Neogastropoda	Harpa	amouretta														1					
Mollusca	Gastropoda	Neogastropoda	Latirolagena	smaragdula																	1		
Mollusca	Gastropoda	Neogastropoda	Latirus	sp1							1												
Mollusca	Gastropoda	Neogastropoda	Mitra	mitra		1																	
Mollusca	Gastropoda	Neogastropoda	Mitra	cf. contracta								1											
Mollusca	Gastropoda	Neogastropoda	Myurella	affinis									1										
Mollusca	Gastropoda	Neogastropoda	Pleuroploca	glabra cf.											1					1			1
Mollusca	Gastropoda	Neogastropoda	Thais	mancinella																			
Mollusca	Gastropoda	Neogastropoda	Turritatus	turritus											1								
Mollusca	Gastropoda	Neogastropoda	Vasum	ceramicum cf.																			
Mollusca	Gastropoda	Nudibranchia	Nembrotha	megalocera																	1		
Mollusca	Gastropoda	Nudibranchia	Phyllidia	varicosa											1						1		1
Mollusca	Gastropoda	Nudibranchia	Phyllidiella	rudmani	1								1										
Mollusca	Gastropoda	Nudibranchia	Phyllidiella	rosans			1						1										
Mollusca	Gastropoda	Trochoidea (super family)	Tectus	maculatus					1														1
Mollusca	Gastropoda	Trochoidea (super family)	Tectus	pyramis																			
Mollusca	Gastropoda	Trochoidea (super family)	Tectus	pyramis																			
Platyhelminthes	Turbellaria	Polycladida	Maiazoon	orsaki																			
Porifera	Calcarea	Clathrinida	Clathrina	sp.																	1		
Porifera	Calcarea	Clathrinida	Leucetta	chagosensis									1										1
Porifera	Calcarea	Clathrinida	Pericarax	sp.																			
Porifera	Demospongiae	Astrophorida	Stelletta	clavosa																1			
Porifera	Demospongiae	Dendroceratida	Dendrylla	sp.																		1	
Porifera	Demospongiae	Dictyoceratida	Carteriospongia	foliascens																			
Porifera	Demospongiae	Dictyoceratida	Dactylospongia	metachromia																			

Table 1 (Con'td)

Phylum	Class	Order	Genera	Species	1	2	3	4	5	6	7	9	10	11	14	17	18	19	21	22	23	24	25	27	28
Porifera	Demospongiae	Dictyoceratida	Hyrtios	<i>erectus</i>	1				1					1										1	1
Porifera	Demospongiae	Dictyoceratida	Lamellosysidea	<i>chlorea</i>																1					
Porifera	Demospongiae	Halicondrida	Auleta	<i>sp.</i>							1														
Porifera	Demospongiae	Halicondrida	Axinella	<i>labyrinthica</i>				1		1	1	1	1	1	1	1									
Porifera	Demospongiae	Halicondrida	Dragnacidon	<i>australe</i>					1														1		
Porifera	Demospongiae	Halicondrida	Liosina	<i>paradoxa</i>				1				1											1		
Porifera	Demospongiae	Halicondrida	Phakellia	<i>carteri</i>	1			1		1	1	1	1	1	1								1		
Porifera	Demospongiae	Halicondrida	Phakellia	<i>cf. cavernosa</i>			1	1	1		1												1		
Porifera	Demospongiae	Halicondrida	Stylissa	<i>flabelliformis</i>			1	1		1	1	1	1	1	1			1							
Porifera	Demospongiae	Haplosclerida	Amphimedon	<i>sp.</i>	1									1			1								
Porifera	Demospongiae	Haplosclerida	Amphimedon	<i>cf. viridis</i>	1		1				1		1			1				1		1			
Porifera	Demospongiae	Haplosclerida	Haliclona	<i>nematifera</i>									1												
Porifera	Demospongiae	Haplosclerida	Haliclona	<i>ostros</i>	1							1					1		1						
Porifera	Demospongiae	Haplosclerida	Microxina	<i>sp.</i>							1				1		1						1		
Porifera	Demospongiae	Haplosclerida	Niphates	<i>sp.</i>															1						
Porifera	Demospongiae	Homoscleromorpha	Plakortis	<i>sp.</i>	1																				
Porifera	Demospongiae	Poecilosclerida	Acamus	<i>sp.</i>	1				1		1			1											
Porifera	Demospongiae	Poecilosclerida	Clathria (Microciona)	<i>plinthina</i>	1					1						1		1							
Porifera	Demospongiae	Poecilosclerida	Monanchora	<i>unguiculata</i>						1						1									
Porifera	Demospongiae	Spirophorida	Craniella	<i>abracadabra</i>												1				1					
Porifera	Demospongiae	Spirophorida	Paratellia	<i>bacca</i>	1		1			1	1	1	1						1				1		
Porifera	Demospongiae	Verongida	Pseudoceratina	<i>verrucosa</i>															1						

STATISTICS


Plate 1: Specimen of the Echinodermata phylum photographed in situ
(photos: Jean-Louis Menou).



Linckia multifora



Choriaster granulatus.



Comanthina schlegeli



Fromia indica



Bohadschia marmorata




Holothuria fuscogilva

Plate 2: Specimen of the Echinodermata phylum photographed in situ
(photos: Jean-Louis Menou).



Culcita cf novaeguineae



Culcita schmedeliana.



Echinostrephus molaris



Echinometra mathaei



Ophiarachnella cf. gorgonia



Ophiothrix sp.

Plate 3: Specimen of the Cnidaria phylum photographed in situ
(photos: Jean-Louis Menou).




Heteractis aurora



Entacmea quadricolor



Chironephthya sp.



Dendronephthya sp.




Acanthogorgia sp.




Cirripathes cf. *anguineus*.

Plate 4: Specimen of the Porifera phylum photographed in situ
(photos: Jean-Louis Menou).



Phakellia carterii




Paratetilla bacca




Carteriospongia foliascens.



Clathria (Microciona) plinthina



Haliclona nematifera



Haliclona ostros